

Cancer Statistics, 2020: Report From National Cancer Registry Programme, India

Prashant Mathur, DNB, PhD¹; Krishnan Sathishkumar, MSc¹; Meesha Chaturvedi, MBBS¹; Priyanka Das, B-Level¹; Kondalli Lakshminarayana Sudarshan, MSc¹; Stephen Santhappan, MSc, MPhil¹; Vinodh Nallasamy, MSc, MPhil¹; Anish John, BE¹; Sandeep Narasimhan, MSc¹; and Francis Selvaraj Roselind, MSc¹; on behalf of ICMR-NCDIR-NCRP Investigator Group

PURPOSE The systematic collection of data on cancer is being performed by various population-based cancer registries (PBCRs) and hospital-based cancer registries (HBCRs) across India under the National Cancer Registry Programme–National Centre for Disease Informatics and Research of Indian Council of Medical Research since 1982.

METHODS This study examined the cancer incidence, patterns, trends, projections, and mortality from 28 PBCRs and also the stage at presentation and type of treatment of patients with cancer from 58 HBCRs (N = 667,666) from the pooled analysis for the composite period 2012–2016. Time trends in cancer incidence rate were generated as annual percent change from 16 PBCRs (those with a minimum of 10 years of continuous good data available) using Joinpoint regression.

RESULTS Aizawl district (269.4) and Papumpare district (219.8) had the highest age-adjusted incidence rates among males and females, respectively. The projected number of patients with cancer in India is 1,392,179 for the year 2020, and the common 5 leading sites are breast, lung, mouth, cervix uteri, and tongue. Trends in cancer incidence rate showed an increase in all sites of cancer in both sexes and were high in Kamrup urban (annual percent change, 3.8%; $P < .05$). The majority of the patients with cancer were diagnosed at the locally advanced stage for breast (57.0%), cervix uteri (60.0%), head and neck (66.6%), and stomach (50.8%) cancer, whereas in lung cancer, distant metastasis was predominant among males (44.0%) and females (47.6%).

CONCLUSION This study provides a framework for assessing the status and trends of cancer in India. It shall guide appropriate support for action to strengthen efforts to improve cancer prevention and control to achieve the national noncommunicable disease targets and the sustainable development goals.

JCO Global Oncol 6:1063-1075. © 2020 by American Society of Clinical Oncology

Creative Commons Attribution Non-Commercial No Derivatives 4.0 License 

INTRODUCTION

Globally, noncommunicable diseases (NCDs) accounted for 71% of total deaths. In India, NCDs were estimated to account for 63% of all deaths, and cancer was one of the leading causes (9%).¹ Cancer registries are recognized as vital components of national cancer-control programs.² Publications from developed and developing countries provide updated information on cancer occurrence, trends, and projections.^{3–7} In India, the systematic collection of data on cancer has been performed since 1982 by the population-based cancer registries (PBCRs) and hospital-based cancer registries (HBCRs) under the National Cancer Registry Programme (NCRP)–National Centre for Disease Informatics and Research (NCDIR) of the Indian Council of Medical Research (ICMR; ICMR-NCDIR-NCRP), Bengaluru (Appendix). NCRP commenced with the objective of generating reliable data on the magnitude and

patterns of cancer. Several NCRP reports on cancer from different registries across India have been published.^{8–9}

PBCRs provide statistics on the occurrence and outcome of cancer in a geographically defined population. They also provide the framework for assessing the control of cancer in the community. HBCRs are concerned with the recording of information on patients with cancer seen in a particular hospital and are mainly used for reviewing clinical performance and the hospital cancer program.¹⁰

This article reports the cancer incidence, patterns, time trends, and mortality from 28 PBCRs for the composite period 2012–2016 across India under ICMR-NCDIR-NCRP. Also, it covers the stage at presentation and type of treatment received by patients with cancer from 58 HBCRs for the period 2012–2016 in India under the

ASSOCIATED CONTENT

Appendix

Data Supplement

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on June 3, 2020 and published at ascopubs.org/journal/go on July 16, 2020; DOI <https://doi.org/10.1200/GO.20.00122>

CONTEXT

Key Objective

This study reports the cancer incidence, patterns, trends, projections, cancer mortality, and clinical aspects of stage at presentation and treatment for the period 2012-2016 from 28 population-based and 58 hospital-based cancer registries in India under the network of the National Cancer Registry Programme.

Knowledge Generated

There is heterogeneity in cancer incidence (age-adjusted rate per 100,000: males: Osmanabad and Beed, 39.5 v Aizawl district, 269.4; females: Osmanabad and Beed, 49.4 v Papumpare district, 219.8) and in patterns across India and a lower proportion (< 33%) of early stage at presentation for common cancers. The projected incidence of patients with cancer for the year 2020 in India in males is 679,421 and in females is 712,758.

Relevance

Results of this study will help in assessing the status and trends of cancer in India. This will assist local- and national-level stakeholders to implement public health action to control cancer.

network of NCRP. Based on estimates from this period, the projection of patients with cancer in India for the year 2020 is also included.

METHODS

Presently, there are 36 PBCRs and 236 HBCRs registered under the ICMR-NCDIR-NCRP. However, this article includes data from 28 PBCRs and 58 HBCRs, which were complete with at least 1 year of good-quality data. All neoplasms with a behavior code of 3 as defined by the International Classification of Diseases for Oncology, 3rd Edition, and the International Statistical Classification of Diseases and Related Health Problems (10th revision; ICD-10) were considered reportable and therefore registered in NCRP.¹¹ Cancer registration is a complex process. In India, cancer registration is active wherein trained registry staff go to different sources (hospitals, diagnostic laboratories, vital statistics departments) for collection of data on a standardized core form.¹⁰ This is followed by quality control checks, duplicate checks, matching with mortality cases, follow-up of death certificate notifications, and assigning death certificate only (DCO). Patients with cancer who were residents in the registration area for a minimum period of 1 year before the date of diagnosis were included in the registry. Multiple cancer data sources were followed for data collection.¹⁰ Quality of the data was maintained per International Association of Cancer Registries/International Agency for Research on Cancer (IACR/IARC) norms.^{12,13}

Incidence and mortality data were retrieved from 28 PBCRs (urban or rural, or both) for the period 2012-2016. Cancer burden measures were calculated as crude rate, age-adjusted rate (AAR) per 100,000 population using world standard population,¹⁴ and cumulative risk (probability that an individual will be diagnosed with cancer [0- to 74-year-old age group] in the absence of any competing cause of death and assuming that the current trends prevail over the time period). Time trends in cancer incidence rate were

generated as annual percent change (APC) and considered statistically significant (at $P < .05$) from 16 PBCRs (with minimum of 10 years of data) using the Joinpoint regression program, 4.0.1 (National Cancer Institute).¹⁵ The years of data for trend analysis varied (11 to 35 years) across the PBCRs, and there were 6 PBCRs with more than 25 years of continuous data.

The country was categorized into 6 geographic zones based on the location of the PBCRs: North (Delhi, Patiala), South (Hyderabad, Kollam, Thiruvananthapuram, Bangalore, and Chennai), East (Kolkata), West (Ahmedabad urban, Aurangabad, Osmanabad and Beed, Barshi rural, Mumbai, and Pune), Central (Wardha, Bhopal, and Nagpur) and Northeast (NE; Manipur, Mizoram, Sikkim, Tripura, West Arunachal, Meghalaya, Nagaland, Pasighat, Cachar, Dibrugarh, and Kamrup urban). Population denominators were derived from the Census of India, conducted by the Registrar General and Census Commissioner of India under the Ministry of Home Affairs, Government of India.¹⁶ The census populations of 2001 and 2011 were used to calculate the postcensal population estimates for the years 2012 and 2016 by 5-year age groups and sex. For time trend analysis, the respective 1981, 1991, 2001, and 2011 censuses were taken as the base to estimate inter- or postcensal population by 5-year age groups and sex.¹⁷

A standardized patient information form for HBCRs under NCRP and the data collection methods have been explained previously.¹⁸ Newly diagnosed and treated patients with cancer (N = 667,666) from 58 HBCRs (with a minimum of 1 year of a complete dataset) in 2012-2016 were pooled and are presented as the relative proportion of clinical stage and treatment. The stage/clinical extent of disease before treatment (excludes previously treated patients with cancer) was classified into localized only, locoregional, distant metastasis, and unknown.¹⁹ The completion of planned cancer-directed treatment after

TABLE 1. Annual Average No. of Patients for All Sites of Cancer With Incidence Rate per 100,000, Cumulative Risk (0-74 age group) and Mortality Rate by Sex and Reporting Year for 28 PBCRs Under NCRP

Serial No.	Registry (period of reference)	Males					Females				
		Incidence			Cum Risk	Mortality	Incidence			Cum Risk	Mortality
		No.	CR	AAR			No.	CR	AAR		
North											
1	Delhi (2012-14)	10,344	112.3	147.0	1 in 6	22.2	9,688	119.6	141.0	1 in 7	17.8
2	Patiala district (2012-16)	1,079	101.6	108.2	1 in 9	32.7	1,215	127.7	124.6	1 in 8	30.1
South											
3	Hyderabad district (2014-16)	1,714	84.2	101.6	1 in 9	15.5	2,151	109.8	136.0	1 in 7	12.5
4	Kollam district (2012-16)	1,986	159.4	127.7	1 in 7	66.5	1,956	139.1	107.1	1 in 9	38.3
5	Thiruvananthapuram district (2012-16)	2,701	170.4	137.8	1 in 7	57.7	2,865	164.8	127.3	1 in 8	39.5
6	Bangalore (2012-14)	4,407	96.8	122.1	1 in 7	42.6	5,276	125.1	146.8	1 in 6	41.5
7	Chennai (2012-16)	2,894	121.8	119.9	1 in 8	35.7	3,361	141.4	132.8	1 in 7	28.8
East											
8	Kolkata (2012-15)	2,547	109.9	91.2	1 in 10	37.9	2,288	105.9	89.2	1 in 11	32.1
West											
9	Ahmedabad urban (2012-16)	2,916	89.1	98.3	1 in 9	27.0	2,205	74.7	76.7	1 in 12	16.9
10	Aurangabad (2012-16)	385	56.6	70.9	1 in 13	13.5	400	62.9	75.1	1 in 12	8.5
11	Osmanabad and Beed (2012-15)	909	39.3	39.5	1 in 23	10.3	1,117	52.8	49.4	1 in 19	10.4
12	Barshi Rural (2012-16)	145	53.9	50.6	1 in 17	35.0	163	67.2	61.0	1 in 15	36.1
13	Mumbai (2012-15)	6,564	97.3	108.4	1 in 9	66.0	6,865	117.6	116.2	1 in 8	61.4
14	Pune (2012-16)	1,937	67.5	83.0	1 in 11	35.3	2,164	83.3	94.0	1 in 10	35.3
Central											
15	Wardha district (2012-16)	478	70.4	64.5	1 in 14	42.3	507	78.7	69.9	1 in 14	37.1
16	Bhopal (2012-15)	892	83.3	101.0	1 in 9	38.3	897	90.4	106.9	1 in 8	30.9
17	Nagpur (2012-16)	1,190	89.0	91.1	1 in 10	21.3	1,209	93.1	89.8	1 in 11	17.7
Northeast											
18	Manipur state (2012-16)	740	47.0	62.8	1 in 14	20.5	900	57.8	71.1	1 in 12	17.3
	Imphal West District (2012-16)	227	85.1	95.3	1 in 9	29.6	300	107.9	110.9	1 in 8	24.3
19	Mizoram state (2012-16)	865	146.1	207.0	1 in 5	121.4	747	127.5	172.3	1 in 5	76.4
	Aizawl district (2012-16)	436	206.2	269.4	1 in 4	152.7	380	174.6	214.1	1 in 5	89.5
20	Sikkim state (2012-16)	234	69.9	88.7	1 in 10	46.4	226	75.3	97.0	1 in 10	46.2
21	Tripura state (2012-16)	1,312	67.0	80.9	1 in 11	46.0	983	52.0	58.3	1 in 16	28.9
22	West Arunachal (2012-16)	244	56.6	101.1	1 in 8	27.3	234	56.3	96.3	1 in 10	18.9
	Papumpare district (2012-16)	94	94.8	201.2	1 in 4	56.5	106	105.1	219.8	1 in 4	37.9
23	Meghalaya (2012-16)	938	92.6	176.8	1 in 5	71.7	566	55.7	96.5	1 in 9	38.1
	East Khasi Hills (2012-16)	577	131.0	227.9	1 in 4	95.0	346	76.9	118.6	1 in 8	51.5
24	Nagaland (2012-16)	281	74.5	124.5	1 in 7	27.8	198	56.3	88.2	1 in 10	11.1
25	Pasighat (2012-16)	64	90.7	120.4	1 in 7	30.9	61	88.1	116.2	1 in 8	22.0
26	Cachar district (2012-16)	933	99.2	129.0	1 in 7	25.2	789	87.0	104.8	1 in 9	17.5
27	Dibrugarh district (2012-16)	507	72.5	91.9	1 in 9	24.0	448	66.0	76.8	1 in 12	14.1
28	Kamrup urban (2012-16)	1,245	190.5	213.0	1 in 4	66.7	958	150.8	169.6	1 in 6	37.3

NOTE. Reporting year data given in parentheses; Meghalaya covers East Khasi Hills, West Khasi Hills, Jaintia Hills, and Ri Bhoi districts; Nagaland covers Kohima and Dimapur districts; Pasighat covers East Siang and Upper Siang; West Arunachal covers Tawang, West Kameng, East Kameng, Upper Subansiri, Lower Subansiri, Kurng Kumey, Papumpare, and West Siang districts.

Abbreviations: AAMR, age-adjusted mortality rate per 100,000; AAR, age-adjusted rate per 100,000; CR, crude rate per 100,000; cum risk, cumulative risk of developing any site of cancer 0-74 years of age; NCRP, National Cancer Registry Program; PBCR, population-based cancer registry.




FIG 1. Comparison of all cancer sites' age-adjusted incidence rates (AARs) of all population-based cancer registries, 2012-2016 (International Statistical Classification of Diseases and Related Health Problems, 10th revision: C00-C97). AARs are in blue and crude rates are given in parentheses in red. Thi'puram district, Thiruvananthapuram district.

diagnosis was classified as surgery, radiotherapy, systemic therapy, and multimodality (combination of surgery and/or radiotherapy and/or systemic therapy).

Good-quality data indices of microscopic verification (MV %) above 75%, DCO% below 20%, other and unspecified sites (O&U%) below 15%, and mortality-to-incidence ratio (M:I%) were calculated for each PBCR and accordingly classified.^{12,13} NCRP has developed in-house software (PBCR and HBCR Data Management) for data capture, quality checks, duplicates checks (deterministic and phonetic-similar sounding duplicate names), and mortality-incidence matching. The list of errors was sent back to registries for clarifications and corrected at each level. Incidence data for 2012-2016 was used as a reference for projection of patients with cancer in India until 2020 by sex and anatomic site (see Data Supplement for more details).

RESULTS

Average annual number of patients for all sites of cancer with incidence rate, cumulative risk by sex, and mortality rate according to different regions in India between 2012 and 2016 are given in Table 1. PBCRs in the NE showed the highest incidence rate in both sexes. It was observed

that Aizawl district had the highest AAR (269.4) and mortality (152.7) rate among males. One of every 4 males in Aizawl district, Papumpare district, East Khasi Hills district, and Kamrup urban are likely to develop cancer in the age group of 0-74 years. One of every 4 females in the Papumpare district and 1 of 5 females in Mizoram state are likely to develop cancer in the age group of 0-74 years.

On comparison of AARs for all sites of cancer (ICD10: C00-C97) across the population among males, Aizawl district (269.4) had the highest AAR and was approximately twice the AAR of Delhi PBCR (147.0). East Khasi Hills district of Meghalaya (227.9) had the second highest AAR, followed by Kamrup urban (213.0) and Mizoram PBCR (207.0). Among females, Papumpare district (219.8) had the highest AAR. In the rest of the regions (excluding NE), Delhi had highest AAR (147.0), followed by Thiruvananthapuram district (137.8) among males; Bangalore (146.8) had highest AAR, followed by Delhi (141.0) among females (Fig 1).

East Khasi Hills district of Meghalaya had the highest relative proportion of cancers associated with the use of tobacco, with 70.4% and 46.5% for males and females,




FIG 2. Sites of cancer associated with the use of tobacco (%), 2012-2016. Crude rate in red and age-adjusted rates in blue are given in parentheses. Sites of cancer associated with the use of tobacco, anatomic sites (International Statistical Classification of Diseases and Related Health Problems, 10th revision codes), lip (C00); tongue (C01-C02); mouth (C03-C06); pharynx (C10 and C12-C14); esophagus (C15); larynx (C32); lung (C33-C34); urinary bladder (C67).³³ Thi'puram district, Thiruvananthapuram district.

respectively. The higher proportion of cancers associated with use of tobacco was in the NE states, followed by registries in the West and Central regions (Fig 2).

Among males, lung, mouth, esophagus, and stomach were the most common cancer sites. Among females, breast cancer, followed by cervix uteri and ovary cancer, were the

most common sites across the PBCRs. Thyroid cancer was the second most common cancer in the PBCRs of Thiruvananthapuram and Kollam, whereas lung cancer was seen in Manipur and Mizoram state. In the NE region, the third most common cancers were stomach and gallbladder (Data Supplement). The decadal changes in leading sites of




FIG 3. Relative proportion (%) of patients according to clinical extent of disease, 2012-2016 (proportion [%] may not total 100% because of rounding).

cancer from 6 older PBCRs (Barshi rural, Bangalore, Bhopal, Chennai, Delhi, and Mumbai) were observed for the first 10 and last 10 years of data (Data Supplement).

The relative proportion of patients according to clinical extent of disease at the time of diagnosis as seen in the pooled data of 58 HBCRs for common sites of cancer showed that the majority of patients with cancer were diagnosed as locally advanced/locoregional for breast (57.0%), cervix uteri (60.0%), head and neck (66.6%), and stomach (50.8%) cancer. The majority of patients with lung cancer were diagnosed with distant metastasis in males (44.0%) and females (47.6%; Fig 3). The relative proportion of types of cancer-directed treatment received (only at the reporting hospital) according to clinical extent of disease before treatment showed that multimodality was the first choice of treatment (locoregional, 79.5%; localized, 74.4%; distant metastasis, 47.6%; Data Supplement).

The estimated APC in cancer AAR for selected anatomic sites of cancer over the time period showed an increase in the incidence rate of all sites of cancer (12 PBCRs in males and 13 PBCRs in females). There was a significant increase in the incidence rate of breast cancer across all PBCRs over the years, except in Nagpur PBCR. There was a significant

decrease in the incidence rate of cervical cancer in 10 PBCRs, except in Dibrugarh district and Pune. Lung cancer showed a significant increase in 11 PBCRs among females (Fig 4).

Among 28 PBCRs, MV%, ranged between 77% (Patiala) and 96.7% (Hyderabad); DCO% ranged between < 0.05% (Osmanabad and Beed) and 19.4% (Patiala); O&U% ranged between 1.8% (Hyderabad) and 13.0% (Patiala). M:I% was high in Barshi rural (67.2%) followed by Wardha (59.2%) and Mumbai (56.0%; Data Supplement).

The projected incidence of patients with cancer in India among males was 679,421 (94.1 per 100,000) and among females 712,758 (103.6 per 100,000) for the year 2020. One in 68 males (lung cancer), 1 in 29 females (breast cancer), and 1 in 9 Indians will develop cancer during their lifetime (0-74 years of age; Table 2). The projected 5 most common cancers in 2020 for males (lung, mouth, prostate, tongue, and stomach) constitute 36% of all cancers and for females (breast, cervix uteri, ovary, corpus uteri, and lung) constitute 53% of all cancers (Data Supplement).

DISCUSSION

India exhibits heterogeneity in cancer. The incidence rates of Aizawl district were observed to be 7 times and 4 times




FIG 4. Annual percentage change (APC) in age-adjusted incidence rates (AAR) over the time period. Calendar years of incidence data for each population-based cancer registry (PBCR) used in trend analysis: Chennai (1982-2016); Mumbai (1982-2015); Bangalore (1982-2014); Barshi rural (1988-2016); Bhopal (1988-2015); Delhi (1988-2014); Kamrup urban (2003-2016); Mizoram state (2004-2016); Dibrugarh district, Sikkim state, Imphal West district; Aurangabad; Thiruvananthapuram (Thi'puram) taluk; Nagpur (2005-2016); Kollam district; Pune (2006-2016); PBCRs with small numbers (< 10 patients) per year not analyzed. Increase in APC in red, decrease in APC in green. (*) APC significantly different from zero; $P < .05$. Thi'puram PBCR expanded its coverage to district from the year 2012 onwards. Hence, Thi'puram taluk data was used for trend analysis. Thi'puram taluk, Thiruvananthapuram taluk.

that of Osmanabad and Beed district PBCRs in males and females, respectively. The highest cancer incidence rate was observed in the NE region (6 PBCRs for males and 4 PBCRs for females) than other areas in the country. The leading sites of cancer in the NE region were nasopharynx, hypopharynx, esophagus, stomach, liver, gallbladder, larynx, lung, breast, and cervix uteri. The NE region lacks required infrastructure with respect to specialized treatment facilities, human resources, as seen by the low 5-year survival of breast, cervix, and head and neck cancer compared with rest of India. A substantial proportion of patients with cancer from the NE region are traveling outside the NE for treatment and cancer care.^{20,21} Local cultural factors and lifestyle choices may have contributed to the heterogeneity in cancer incidence pattern and differences in India, as was seen in Thailand.⁷

Lung (9 PBCRs), mouth (9 PBCRs), esophagus (5 PBCRs), stomach (4 PBCRs), and nasopharynx (1 PBCR) cancers were the most common cancers in men. Lung cancer was the leading site in metropolitan cities and the southern region, whereas mouth cancer was the leading site in the West and Central regions. Lung cancer and oral/mouth cancer were the most common cancers among males in the Indian subcontinent.²² Cancers of the esophagus, stomach, and nasopharynx were the leading sites in the NE region of India. Here, the cancer incidence pattern is different from the rest of India. There are similarities in the

cancer incidence pattern with the Southeast Asian region.²³ Overall, these findings on patterns of cancer were similar to previously published reports under NCRP.⁸

Cancer of the breast (19 PBCRs) and cervix uteri (7 PBCRs) were the most common cancers in women. The highest burden of breast cancer was observed in metropolitan cities. There is an increase in the trend of incidence of breast cancer, whereas cervix uteri cancer is on the decline. A steady increase in breast cancer in most of the PBCRs including newer PBCRs, poses a great health challenge to women in India.²⁴ Presently, breast cancer and cervix uteri are the leading sites of cancer among women in India, posing an important public health problem that needs important input from various health and other agencies to tackle.²⁵ A multidisciplinary approach to breast cancer, including awareness programs, preventive measures, screening programs for early detection, and availability of treatment facilities, are vital for reducing both incidence and mortality of cancer in Indian women.²⁶

The incidence rate of thyroid cancer among women is increasing, and it is most common in the districts of Thiruvananthapuram and Kollam in Kerala. The high burden of thyroid cancer in Kerala could be due to overdiagnosis,²⁷ as was observed even in high-income and low- and middle-income countries.²⁸ AAR in Barshi rural is almost one third of urban PBCRs (males, 50.6 v 147.0; females, 61.0 v 146.8), and the increase in APC was

TABLE 2. Projected Incidence of Cancer Statistics in India, 2020

Site	Male			Female			Both Sexes		
	Patients	CR	Cum Risk	Patients	CR	Cum Risk	Patients	CR	Cum Risk
All sites	679,421	94.1	1 in 9	712,758	103.6	1 in 9	1,392,179	98.7	1 in 9
Oral cavity and pharynx	139,018	19.2	1 in 41	49,951	7.3	1 in 112	188,969	13.4	1 in 60
Tongue	39,902	5.5	1 in 147	13,870	2.0	1 in 401	53,772	3.8	1 in 215
Mouth	57,380	7.9	1 in 103	22,483	3.3	1 in 241	79,863	5.7	1 in 144
Pharynx	3,029	0.4	1 in 1,793	1,102	0.2	1 in 5,475	4,131	0.3	1 in 2,701
Other oral cavity	38,707	5.4	1 in 137	12,496	1.8	1 in 476	51,203	3.6	1 in 213
Digestive system	163,845	22.7	1 in 32	110,137	16.0	1 in 50	273,982	19.4	1 in 39
Esophagus	32,622	4.5	1 in 159	20,206	2.9	1 in 264	52,828	3.7	1 in 198
Stomach	32,713	4.5	1 in 160	17,430	2.5	1 in 319	50,143	3.6	1 in 213
Small intestine	2,155	0.3	1 in 2,492	1,451	0.2	1 in 3,901	3,606	0.3	1 in 3,044
Colon	20,572	2.8	1 in 260	15,685	2.3	1 in 348	36,257	2.6	1 in 298
Rectum	21,915	3.0	1 in 244	14,985	2.2	1 in 372	36,900	2.6	1 in 295
Anus, anal canal	2,897	0.4	1 in 1,865	2,028	0.3	1 in 2,682	4,925	0.3	1 in 2,200
Liver and intrahepatic bile duct	26,678	3.7	1 in 189	10,732	1.6	1 in 514	37,410	2.7	1 in 277
Gallbladder and other biliary	12,385	1.7	1 in 422	19,510	2.8	1 in 284	31,895	2.3	1 in 340
Pancreas	11,908	1.6	1 in 429	8,110	1.2	1 in 657	20,018	1.4	1 in 519
Respiratory system	103,552	14.3	1 in 48	32,480	4.7	1 in 165	136,032	9.6	1 in 74
Larynx	27,146	3.8	1 in 184	3,316	0.5	1 in 1,633	30,462	2.2	1 in 331
Lung and bronchus	71,788	9.9	1 in 68	26,490	3.9	1 in 201	98,278	7.0	1 in 101
Other respiratory organs	4,618	0.6	1 in 1,273	2,674	0.4	1 in 2,156	7,292	0.5	1 in 1,600
Bones and joints	8,115	1.1	1 in 1,013	5,840	0.8	1 in 1,370	13,955	1.0	1 in 1,162
Soft tissue	8,047	1.1	1 in 842	6,590	1.0	1 in 1,052	14,637	1.0	1 in 936
Skin (excluding basal and squamous)	11,203	1.6	1 in 510	8,962	1.3	1 in 640	20,165	1.4	1 in 568
Melanoma of the skin	3,003	0.4	1 in 1,904	2,364	0.3	1 in 2,281	5,367	0.4	1 in 2,075
Other nonepithelial skin	8,200	1.1	1 in 695	6,598	1.0	1 in 890	14,798	1.0	1 in 781
Breast	5,377	0.7	1 in 1,022	205,424	29.9	1 in 29	210,801	15.0	1 in 56
Genital system	51,994	7.2	1 in 105	155,630	22.6	1 in 36	207,624	14.7	1 in 54
Uterine cervix	—	—	—	75,209	10.9	1 in 75	75,209	10.9	1 in 75
Uterine corpus	—	—	—	26,514	3.9	1 in 190	26,514	3.9	1 in 190
Ovary	—	—	—	43,886	6.4	1 in 133	43,886	6.4	1 in 133
Vulva	—	—	—	2,138	0.3	1 in 2,459	2,138	0.3	1 in 2,459
Vagina and other genital, female	—	—	—	7,570	1.1	1 in 745	7,570	1.1	1 in 745
Placenta	—	—	—	313	0.0	1 in 30,912	313	0.0	1 in 30,912
Prostate	41,532	5.7	1 in 125	—	—	—	41,532	5.7	1 in 125
Testis	4,352	0.6	1 in 2,095	—	—	—	4,352	0.6	1 in 2,095
Penis and other genital, male	6,110	0.8	1 in 916	—	—	—	6,110	0.8	1 in 916
Urinary system	33,269	4.6	1 in 158	11,265	1.6	1 in 502	44,534	3.2	1 in 240
Urinary bladder	20,470	2.8	1 in 250	5,403	0.8	1 in 1,014	25,873	1.8	1 in 402
Kidney and renal pelvis	12,363	1.7	1 in 442	5,657	0.8	1 in 1,038	18,020	1.3	1 in 620

(Continued on following page)

TABLE 2. Projected Incidence of Cancer Statistics in India, 2020 (Continued)

Site	Male			Female			Both Sexes		
	Patients	CR	Cum Risk	Patients	CR	Cum Risk	Patients	CR	Cum Risk
Ureter and other urinary organs	436	0.1	1 in 10,843	205	0.0	1 in 21,892	641	0.0	1 in 14,531
Eye and orbit	1,304	0.2	1 in 6,870	953	0.1	1 in 9,063	2,257	0.2	1 in 7,792
Brain and other nervous system	19,979	2.8	1 in 341	12,750	1.9	1 in 546	32,729	2.3	1 in 419
Endocrine system	9,263	1.3	1 in 709	26,665	3.9	1 in 279	35,928	2.5	1 in 402
Thyroid	8,570	1.2	1 in 759	26,095	3.8	1 in 285	34,665	2.5	1 in 416
Adrenal gland	693	0.1	1 in 10,797	570	0.1	1 in 14,053	1,263	0.1	1 in 12,209
Lymphoma	32,695	4.5	1 in 197	20,247	2.9	1 in 296	52,942	3.8	1 in 236
Hodgkin lymphoma	7,294	1.0	1 in 1,150	3,936	0.6	1 in 1,871	11,230	0.8	1 in 1,418
Non-Hodgkin lymphoma	25,344	3.5	1 in 238	16,263	2.4	1 in 352	41,607	3.0	1 in 284
Malig Imn Prol D	57	0.0	1 in 101,774	48	0.0	1 in 164,355	105	0.0	1 in 126,050
Multiple myeloma	10,725	1.5	1 in 465	7,756	1.1	1 in 646	18,481	1.3	1 in 541
Leukemia	32,481	4.5	1 in 239	21,132	3.1	1 in 353	53,613	3.8	1 in 284
Lymphoid leukemia	14,159	2.0	1 in 608	7,419	1.1	1 in 1,138	21,578	1.5	1 in 789
Myeloid leukemia	14,913	2.1	1 in 474	11,275	1.6	1 in 617	26,188	1.9	1 in 536
Leukemia uns	3,409	0.5	1 in 2,287	2,438	0.4	1 in 2,980	5,847	0.4	1 in 2,583
Other and unspecified primary sites	48,554	6.7	1 in 114	36,976	5.4	1 in 153	85,530	6.1	1 in 131

Abbreviations: CR, crude rate; cum risk, cumulative risk of developing cancer at 0-74 years of age; Malig Imn Prol D, malignant immunoproliferative diseases; Uns, unspecified.

less compared with urban PBCRs. This needs additional investigation.

There are cancers of several anatomic sites known to be associated with the use of tobacco.²⁹ Based on PBCR data, almost one third of the cancers were known to be associated with the use of tobacco in India. India state-level disease burden initiative cancer collaborators estimated that tobacco use was the highest contributing risk factor for cancer in India. In India, lung cancer can be attributed to tobacco use and air pollution, which are the leading risk factors.³⁰ Approximately 70% of cancers in India were potentially preventable through modifiable risk factors.³¹

Because it is difficult to obtain information on the clinical extent of disease and treatment from PBCRs, the hospital database was used for such analysis.¹⁰ The majority of breast and cervix uteri cancers were diagnosed at a locally advanced stage. Chemoradiation was the most common type of treatment of cancer cervix uteri. A multi-institutional study from India on cervix cancer showed significantly better survival with chemoradiation than radiation alone in the locally advanced stage.¹⁸ A study from Chennai showed that concurrent chemoradiation for locally advanced cervical cancer resulted in the best disease-free survival.³² Two thirds of the patients with cancer were diagnosed at the locoregional stage for head and neck cancers from HBCRs. Similar to that, a low proportion of patients with head and neck cancer presented in the early stage, and a high

proportion (88.1%) were seen in Uttarakhand.³³ Multi-modality was the most common treatment given for breast and head and neck cancers. A multi-institutional study estimated that 65% of new head and neck cancers with locally advanced disease did not receive the benefit of optimal treatment, resulting in poor survival.³⁴

Less than one fifth of lung and stomach cancers were diagnosed as localized only. Systemic therapy was the most common type of treatment given for lung and stomach cancer. A previous report on HBCR results showed similar findings.^{8a} A hospital-based study from northern India showed that 90% of patients with lung cancer were diagnosed at an advanced stage of the disease, and there was a delay in diagnostic evaluation and treatment.³⁵ Creating cancer awareness, preventing risk factors, and improving access to care among people would result in downstaging of cancer.

The measure of validity, MV%, was above 77% for all the PBCRs. Varying patterns of DCO% and M:I% were observed among PBCRs which were dependent on the quality of death registration and certification.¹³ Efforts to improve the quality are always underway. In some registries, low DCO% (< 1%) is due to nonavailability of all-cause mortality data and incomplete/incorrect certification of cause of death. Some registries had an efficient trace back procedure by house visit/phone. Data from PBCRs were regularly published in successive volumes of Cancer Incidence

in Five Continents (CI-5) by WHO-IACR/IARC. The incidence data from 15 PBCRs under NCRP (India) were published in Cancer Incidence in Five Continents, Volume XI, by WHO-IACR/IARC.³⁶⁻³⁸

The projected incidence of patients with cancer is higher for females (712,758) than males (679,421) for the year 2020. The projected national cancer incidence burden in 2020 will be 98.7 per 100,000 population (1,392,179 patients) as a conservative estimate. It is assumed that the observed rate of 2012-2016 will remain unchanged until 2020. The time trend in rate was not used to avoid uncertainty in the projection for a populous country like India. NCRP has estimated a slightly higher number of patients with cancer compared with IACR/IARC and GLOBOCAN for all sites of cancer in 2018. This may be because of a difference in methodology and use of recent data (1,392,179 v 1,157,294).²² This is the first such attempt in the country and will be further updated on availability of the next data set and census information. The influencing factors, such as risk factors/behavior, case finding procedure, screening program, and improved techniques for detecting patients

with cancer, are likely to influence the projected number of patients.

PBCRs in this study covered 100 million average annual person-years, accounting for coverage close to 10% of the population in India. Cancer registration in India faces several challenges because it is not a notifiable disease, posing challenges to data collection.³⁹⁻⁴² The mortality registration system has several gaps, including incomplete and inaccurate certification of cause of death.^{43,44} Registering through passive notification by health care providers to report cancer occurrence in India would improve the coverage with limited resources. Linking of cancer registry data with Ayushman Bharat,⁴⁵ mortality databases, and the Hospital Information System would improve cancer registration, follow-up, and outcome data.

This study provides a framework for assessing the status and trends of cancer in India. This shall guide appropriate support for action to strengthen efforts to improve cancer prevention and control to achieve the National NCD targets and the sustainable development goals.^{46,47} The data also provide leads to key research questions.

AFFILIATION

¹National Centre for Disease Informatics and Research, Bengaluru, India

CORRESPONDING AUTHOR

Prashant Mathur, PhD, National Centre for Disease Informatics and Research, ICMR Complex (II Floor), Poojanahalli, Kannamangala Post, Bengaluru, Karnataka 562 110 India; Twitter: @ncdirindia; e-mail: director@ncdirindia.org.

AUTHOR CONTRIBUTIONS

Conception and design: Prashant Mathur, Krishnan Sathishkumar, Meesha Chaturvedi

Collection and assembly of data: Prashant Mathur, Krishnan Sathishkumar, Meesha Chaturvedi, Priyanka Das, Kondalli Lakshminarayana Sudarshan, Anish John, Francis Selvaraj Roselind

Data analysis and interpretation: All authors

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/go/site/misc/authors.html.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](http://OpenPayments)).

No potential conflicts of interest were reported.

ACKNOWLEDGMENT

The authors acknowledge the contribution of the 28 population-based cancer registries and 58 hospital-based cancer registries under the National Cancer Registry Program, which contributed data, and Research Area Panel Expert Group on Cancer.

REFERENCES

1. WHO: World Health Statistics 2019: Monitoring Health for the SDGs. Geneva, Switzerland, World Health Organization, 2018
2. Parkin DM: The evolution of the population-based cancer registry. *Nat Rev Cancer* 6:603-612, 2006
3. Nandakumar A, Gupta PC, Gangadharan P, et al: Geographic pathology revisited: Development of an atlas of cancer in India. *Int J Cancer* 116:740-754, 2005
4. Swaminathan R, Selvakumaran R, Esmey PO, et al: Cancer pattern and survival in a rural district in South India. *Cancer Epidemiol* 33:325-331, 2009
5. Jemal A, Ward EM, Johnson CJ, et al: Annual report to the nation on the status of cancer, 1975-2014, featuring survival. *J Natl Cancer Inst* 109:djx030, 2017
6. de Camargo B, de Oliveira Santos M, Rebelo MS, et al: Cancer incidence among children and adolescents in Brazil: First report of 14 population-based cancer registries. *Int J Cancer* 126:715-720, 2010
7. Pongnikorn D, Daoprasert K, Waisri N, et al: Cancer incidence in northern Thailand: Results from six population-based cancer registries 1993-2012. *Int J Cancer* 142:1767-1775, 2018
8. National Centre for Disease Informatics and Research: Consolidated Report of Population Based Cancer Registries, 2006-2008, 2009-2011, 2012-2014 Bengaluru, India, National Cancer Registry Programme (NCRP-ICMR) <https://ncdirindia.org/Reports.aspx>
- 8a. National Centre for Disease Informatics and Research: Consolidated Report of Hospital Based Cancer Registries, 2004-2006, 2007-2011, 2012-2014 Bengaluru, India, National Cancer Registry Programme (NCRP-ICMR) <https://ncdirindia.org/Reports.aspx>

9. National Centre for Disease Informatics and Research: Time trends in cancer incidence rates, 1982-2010, Bangalore: National Cancer Registry Programme (NCRP-ICMR), 2013. https://www.ncdirindia.org/All_Reports/TREND_REPORT_1982_2010/
10. Jensen OM, Parkin DM, MacLennan R, et al (eds): Cancer Registration: Principles and Methods. Lyon, France, IARC Scientific Publications, 1991
11. WHO: International Statistical Classification of Diseases and Related Health Problems. Geneva, Switzerland, World Health Organization, 1994
12. Bray F, Parkin DM: Evaluation of data quality in the cancer registry: Principles and methods. Part I: Comparability, validity and timeliness. *Eur J Cancer* 45:747-755, 2009
13. Parkin DM, Bray F: Evaluation of data quality in the cancer registry: Principles and methods Part II. Completeness. *Eur J Cancer* 45:756-764, 2009
14. Segi M: Cancer Mortality for Selected Sites in 24 Countries 1950-1957. Sendai, Japan, The Department of Public Health, Tohoku University School of Medicine 1960
15. Joinpoint regression program, version 4.0.1. Statistical research and application branch, National Cancer Institute, 2013. <https://surveillance.cancer.gov/joinpoint/>
16. Census of India: Registrar General of India, Socio Cultural Tables, C14, Population by Five Year Age Group, by Residence and Sex, New Delhi, 2001 & 2011. www.censusindia.gov.in
17. Takiar R, Shobana B: Cancer incidence rates and the problem of denominators - a new approach in Indian cancer registries. *Asian Pac J Cancer Prev* 10:123-126, 2009
18. Nandakumar A, Kishor Rath G, Chandra Katakai A, et al: Concurrent chemoradiation for cancer of the cervix: Results of a multi-institutional study from the setting of a developing country (India). *J Glob Oncol* 1:11-22, 2015
19. Edge SB: AJCC Cancer Staging Manual (ed 7). New York, NY, Springer, 2010
20. National Centre for Disease Informatics and Research: Report on Cancer Burden in North Eastern States of India. Bengaluru, India, National Cancer Registry Programme (NCRP-ICMR) <http://ncdirindia.org/>
21. Ngaihte P, Zomawia E, Kaushik I: Cancer in the NorthEast India: Where we are and what needs to be done? *Indian J Public Health* 63:251-253, 2019
22. Bray F, Ferlay J, Soerjomataram I, et al: Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 68:394-424, 2018
23. Sharma JD, Kalit M, Nirmolia T, et al: Cancer: Scenario and relationship of different geographical areas of the globe with special reference to North East-India. *Asian Pac J Cancer Prev* 15:3721-3729, 2014
24. Chaturvedi M, Vaitheeswaran K, Satishkumar K, et al: Time trends in breast cancer among Indian women population: An analysis of population based cancer registry data. *Indian J Surg Oncol* 6:427-434, 2015
25. Takiar R: Status of breast and cervix cancer in selected registries of India. *Ann Womens Health* 2:1012, 2018
26. Malvia S, Bagadi SA, Dubey US, et al: Epidemiology of breast cancer in Indian women. *Asia Pac J Clin Oncol* 13:289-295, 2017
27. Mathew IE, Mathew A: Rising thyroid cancer incidence in Southern India: An epidemic of overdiagnosis? *J Endocr Soc* 1:480-487, 2017
28. Lortet-Tieulent J, Franceschi S, Dal Maso L, et al: Thyroid cancer "epidemic" also occurs in low- and middle-income countries. *Int J Cancer* 144:2082-2087, 2019
29. WHO: Overall Evaluations of Carcinogenicity: An Updating of IARC Monographs, Volumes 1-42. Lyon, France, IARC, 1987
30. India State-Level Disease Burden Initiative Cancer Collaborators: The burden of cancers and their variations across the states of India: The Global Burden of Disease Study 1990-2016. *Lancet Oncol* 19:1289-1306, 2018
31. Gandhi AK, Kumar P, Bhandari M, et al: Burden of preventable cancers in India: Time to strike the cancer epidemic. *J Egypt Natl Canc Inst* 29:11-18, 2017
32. Shanta V, Selvaluxmy G, Swaminathan R, et al: Evolution in the management of locally advanced cervical cancer: The experience of Cancer Institute (WIA), Chennai, India. *Asian Pac J Cancer Prev* 11:1091-1098, 2010
33. Pandey KC, Revannasiddaiah S, Pant NK, et al: Stage-wise presentation of non-metastatic head and neck cancer: An analysis of patients from the Kumaon hills of India. *Asian Pac J Cancer Prev* 15:4957-4961, 2014
34. Nandakumar A, Kishor Rath G, Chandra Katakai A, et al: Survival in head and neck cancers: Results of a multi-institution study. *Asian Pac J Cancer Prev* 17:1745-1754, 2016
35. Chandra S, Mohan A, Guleria R, et al: Delays during the diagnostic evaluation and treatment of lung cancer. *Asian Pac J Cancer Prev* 10:453-456, 2009
36. Curado MP, Edwards B, Shin HR, et al (eds): Cancer Incidence in Five Continents, Volume IX. Lyon, France, IARC Scientific Publications, 2007
37. Forman D, Bray F, Brewster DH, et al (eds): Cancer Incidence in Five Continents Volume X. Lyon, France, IARC Scientific Publications, 2014
38. Bray F, Colombet M, Mery L, et al (eds): Cancer Incidence in Five Continents, Volume XI. Lyon, France, IARC Scientific Publications, 2017
39. Bray F, Znaor A, Cueva P, et al (eds): Planning and Developing Population-Based Cancer Registration in Low- and Middle-Income Settings. Lyon, France, IARC Technical Publication, 2014
40. Dhar M: A critical review of evolution of cancer registration in India. *J Tumor Med Prev* 2:555594 2018
41. Sahoo SS, Verma M, Parija PP: An overview of cancer registration in India: Present status and future challenges. *Oncol J India* 2:86-89, 2018
42. Behera P, Patro BK: Population based cancer registry of India – the challenges and opportunities. *Asian Pac J Cancer Prev* 19:2885-2889, 2018
43. Kakkar R, Aggarwal P: Civil registration & sample registration system: Challenges and initiatives. *SRHU Medical Journal* 1:47-49, 2017
44. ORGI: Report on Medical Certification of Cause of Death (MCCD) 2017, India, Office of the Registrar General, Ministry of Home Affairs. *Vital Stat (Ott Annu Ed)*, 2017. https://censusindia.gov.in/2011-Documents/mccd_Report1/MCCD_Report-2017.pdf
45. National Health Authority: Ayushman Bharat Pradhan Mantri Jan Arogya Yojana. National Health Authority, Government of India. <http://www.pmjay.gov.in/>
46. WHO: NCD Global Monitoring Framework. https://www.who.int/nmh/global_monitoring_framework/2013-11-06-who-dc-c268-whp-gap-ncds-techdoc-def3.pdf?ua=1
47. United Nations: Transforming our world: The 2030 agenda for sustainable development. <https://sustainabledevelopment.un.org/content/documents/21252030%20Agenda%20for%20Sustainable%20Development%20web.pdf>



APPENDIX

INDIAN COUNCIL OF MEDICAL RESEARCH–NATIONAL CENTRE FOR DISEASE INFORMATICS AND RESEARCH–NATIONAL CANCER REGISTRY PROGRAM INVESTIGATOR GROUP

Swetha Acharya, Dillip Kumar Agarwalla, Zarika Ahmed, Gazi Naseem Ahmed, Shiraj Ahmed, B.S. Ajaikumar, P. Anandhi, Anil Kumar Anand, Ramandeep Arora, R.A. Badwe, Manas Ranjan Baisakh, Balamuralikrishna, Debabrata Barmon, Sangita Bhandari, Samir Bhattacharya, M. Bhattacharyya, Arup Bhowmik, T.W. Bhutia, T. Bindhu, Vijay Kumar Bodal, Pradeep Chandrakar, Latha Chaturvedua, Richa Chauhan, Saia Chenkual, S. Chhabra, Lily Chhakchhuak, Vivek Choudhary, Ashok Kumar Das, Rituparna Das, Dhritiman Datta, Biswajit Debbamma, Shiromani Debbarma, S.V. Suryanarayana Deo, Vinay Deshmane, Madhubala Devi, Punyabati Devi, Laishram Purnima Devi, Rajesh Dikshit, H. Dkhar, Biswajit Dubashi, Debanarayan Dutta, Vijay Gadagi, Vinay Gadigi, B. Ganesh, M.S. Ganesh, K.V. Gangadharan, Nitin Gangane, Uma Garg, Preethi Sara George, Sushmita Ghosal, Radha Ghosh, Debasis Gochhait, Ajay Gogia, H.K. Goswami, Ashutosh Gupta, Manish Gupta, Kaveri Hallikeri, Rajesh Harsvardhan, B.H. Srinivas, Vandana Jain, Swarima Jaitley, K.L. Jayakumar, P. Jayalekshmi, Kaling Jerang, G. Jongkey, J. Seios, K. Vijayakumar, D. Kadambari, Geeta Kadayapath, Ravi Kannan, R. Kapoor, Tejinder Kataria, Mohanvir Kaur, Swarn Kaur, K. Gunaseelan, Shehna A. Khader, V. Khamo, Nazir Ahmad Khan, Sushma Khurajam, Marla K. Prashanth, Anjali Kolhe, S.S. Koyande, Adishi Kri, Dinesh Kumar, G. Dilip Kumar, Narender Kumar, S. Shiva Kumar, Sunil Kumar, Rajesh Singh Laishram, Lalchhanhimi, K. Lalfakzuala, Lalhupuii, Doris Lallawmzuali, Jerry Lalrinsanga, Cindy Lalthanpuii, Imliwati Longkumer, M.B. Jayaraman, Gautam Majumdar, Reeni Malik, Swapna Maliye, Indranil Mallick, Syamsundar Mandal, N. Manoharan, Lone Mohammad Maqbool, Aleyamma Mathew, Sanjay Mishra, Janmejya Mohapatra, C.T. Muthukumar, Rekha G. Nair, Prem Nair, Sunithra Nair, Durgaprasad Nanda, D.K. Nath, Partha Nath, N. Venkatesh, Gamba Padu, Rakesh Pandey, Shashank Pandya, Sangeeta Pankaj, N.S. Panse, Sujith Chayu Patanaik, P.C. Pathy, Jeremy L. Pautu, K. Pavithran, Prasanth Penumade, Priya Darshini Pradhan, Aruna E. Prasad, Punitalal, Kedozeto Punyu, K. Kalpana Raghunath, Mujeeb Rahman, Vinod Raina, Ravi Shankar Rajendra, Ramachandra, C. Ramesh, B.S. Ramesh, Rakesh S. Ramesh, R. Rama Ranganathan, Rita Rani, K. Ratheesan, Sudhir Rawal, Prarthana Roselil, G. Sadashivudu, B.J. Saikia, Projnan Saikia, S.K. Samantara, A. Santa, Anupam Sarma, N. Hariharanadha Sarma, Sunil Kumar Saxena, Pooja Sethi, Anand Shah, B.K. Sharma, Jagannath Dev Sharma, Sudarshan Sharma, Shah Alam Sheikh, Kamalaksh Shenoy, Dinesh Shet, Atul Shrivastava, B.R. Shrivastava, Sushma Shrivastava, Alben Sigamani, Neetu Singhal, Indibor Singh, Rajesh Kumar Singh, Usha Singh, Badarisha R. Sohliya, G.S. Sreenath, U. Srihari, V.G. Sudhakaran, Saina Sunilkumar, Partha Sarathi Sutradhar, R. Swaminathan, A Talukdar, Ritesh Tapkire, Sopai Tawsik, B. Paul Thaliath, K.T. Harichandrakumar, Pampa Ch. Toi, Shyam Tsering, B.B. Tyagi, Ajith Kumar, V.R. Ashok Kumar Vaid, K. Velavan, C.R. Vijay, W.B. Langstieh, Chaitali Wagmere, Meenu Walia, John Zohmingthanga, Eric Zomawia, and B. Zothankima.

Names Listed Alphabetically

Affiliations. National Centre for Disease Informatics and Research, Indian Council of Medical Research, Bengaluru, India: Prashant Mathur, DNB, PhD; Krishnan Sathishkumar, MSc; Meesha Chaturvedi, MBBS; Priyanka Das, B-Level; Kondalli Lakshminarayana Sudarshan, MSc; Stephen Santhappan, MSc, MPhil; Vinodh Nallasamy, MSc, MPhil; Anish John, BE; Sandeep Narasimhan, MSc; and Francis Selvaraj Roselind, MSc.

Population-based cancer registries. Delhi population-based cancer registry (PBCR) – B.R. Ambedkar Institute Rotary Cancer Hospital, All India Institute of Medical Sciences, New Delhi (S.V.

Suryanarayana Deo, MS, N. Manoharan); Patiala District PBCR – Government Medical College and Rajindra Hospital, Punjab (Vijay Kumar Bodal, MD, Mohanvir Kaur, MD); Hyderabad PBCR – Nizam's Institute of Medical Sciences, Telangana (G. Sadashivudu, MD, U. Srihari); Kollam PBCR – District Cancer Registry Kollam, Regional Cancer Centre, Kerala (Rekha G Nair, MD, P. Jayalekshmi, PhD); Thiruvananthapuram PBCR – Regional Cancer Centre, Thiruvananthapuram, Kerala (Aleyamma Mathew, PhD, Preethi Sara George, PhD); Bengaluru PBCR – Kidwai Memorial Institute of Oncology, Karnataka (Ramachandra, MS, C. Ramesh, PhD, C.R. Vijay, PhD); Chennai PBCR – Cancer Institute (WIA), Tamil Nadu (R. Swaminathan, PhD, R. Rama Ranganathan, PhD); Kolkata PBCR – Chittaranjan National Cancer Institute and Saroj Gupta Cancer Centre and Research Institute, West Bengal (Samir Bhattacharya, PhD, Syamsundar Mandal, PhD, Durgaprasad Nanda, MS); Ahmedabad urban PBCR – Gujarat Cancer & Research Institute, Gujarat (Shashak Pandya, MS, Anand Shah, MD); Mumbai, Aurangabad, Nagpur, Pune PBCRs – Indian Cancer Society, Maharashtra (Vinay Deshmane, MD, S.S. Koyande); Barshi Expanded (Osmanabad and Beed Districts) PBCR – Nargis Dutt Memorial Cancer Hospital, Maharashtra (NS Panse); Barshi Rural PBCR – Tata Memorial Hospital, Maharashtra (R.A. Badwe, MS, Rajesh Dikshit, PhD); Wardha PBCR – Mahatma Gandhi Institute of Medical Sciences, Maharashtra (Nitin Gangane, MD, Swapna Maliye, S. Chhabra, MD); Bhopal PBCR – Gandhi Medical College, Madhya Pradesh (Reeni Malik, MD, Atul Shrivastava); Manipur PBCR – Regional Institute of Medical Sciences, Manipur (Madhubala Devi, MD, Punyabati Devi, MD, Laishram Rajesh, MD, Sushma Khurajam); Mizoram PBCR – Civil Hospital, Mizoram (Eric Zomawia, MD, John Zohmingthanga, MD, Lalchhanhimi, MD, Saia Chenkual, MD, B. Zothankima, MD, Lily Chhakchhuak, MBBS, Jerry Lalrinsanga, MBBS); Sikkim PBCR – Sir Thutob Namgyal Memorial Referral Hospital, Sikkim (T.W. Bhutia, MD, Priya Darshini Pradhan, MD, Sangita Bhandari); Naharlagun (West Arunachal) PBCR – Timo Riba State Hospital, Arunachal Pradesh (Sopai Tawsik, MD, Shyam Tsering, MD, K.R. Adishi, MD, Gamba Padu, MD); Meghalaya PBCR – Civil Hospital, Meghalaya (W.B. Langstieh, H. Dkha, MD, B. Sohliya, MD, L Purnima Devi, MD); Nagaland PBCR – Naga Hospital Authority, Nagaland (V. Khamo, MD, Kedozeto Punyu, MD); Pasighat PBCR – Bakin Pertin General Hospital, Arunachal Pradesh (Kaling Jerang, PGDCP, G. Jongkey, MD); Tripura PBCR – Cancer Hospital, Regional Cancer Centre, Tripura (Gautam Majumdar, MD, Arup Bhowmik, MD, Rituparna Das, MD, Shiromani Debbarma); Cachar PBCR – Silchar Medical College, Assam (Shah Alam Sheikh, MD, R.P. Banik, MD); Dibrugarh PBCR – Assam Medical College and Hospital, Assam (Projnan Saikia, MD, Zarika Ahmed, MD); Guwahati (Kamrup urban) PBCR – Dr. Bhubaneswar Borooah Cancer Institute (Regional Institute for Treatment and Research), Assam (Jagannath D. Sharma, MD, Anupam Sarma, MD, Debabrata Barmon, MD, Shiraj Ahmed, MD).

Hospital-Based Cancer Registries A.J. Hospital & Research Centre, Mangalore (Marla K. Prashanth, MS, Kamalaksh Shenoy, MD); Acharya Harihar Regional Cancer Centre, Cuttack (Dillip Kumar Agarwalla, MD, Janmejya Mohapatra, MD, P.C. Pathy, MD, S K Samantara, MD); Amrita Institute of Medical Sciences and Research, Kochi (Debanarayan Dutta, MD, K. Vijayakumar, MS, Prem Nair, MD, K. Pavithran, MD); Apollo Hospital, Bhubaneswar (Manas Ranjan Baisakh, MD); Asian Institute of Medical Sciences, Faridabad (Neetu Singhal, DNB); Assam Medical College, Dibrugarh (H. K. Goswami, MD); BPS Government Medical College for Women, Sonipat (Uma Garg, MD, Swarn Kaur, MD); Cachar Cancer Hospital, Silchar (Ravi Kannan, MS, Ritesh Tapkire, MS), Cancer Hospital & Research Institute, Gwalior (B.R. Shrivastava, MS); Cancer Institute (WIA), Chennai (R. Rama Ranganathan, PhD, R. Swaminathan, PhD); Chittaranjan National Cancer Institute, Kolkata (Partha Nath, MD, Syamsundar Mandal, PhD); B. Borooah Cancer Institute, Guwahati (M Bhattacharyya, MD, Ashok Kumar Das, MS, B J. Saikia, MD, A Talukdar, MS, Jagannath Dev Sharma, MD); B.R. Ambedkar Institute Rotary Cancer Hospital, New Delhi (Ajay Gogia, MS, Sunil Kumar, MS, S.V. Suryanarayana Deo, MS, N. Manoharan, G.K. Rath, MD); Erode Cancer Centre, Thindal, Erode (K. Velavan, MD); Father Muller Medical College Hospital, Mangalore (Dinesh Shet, MD); Fortis Memorial

Research Institute, Gurugaon (Vinod Raina, MD, B. B. Tyagi, MS); Gandhi Medical College, Bhopal (Atul Shrivastava, Sushma Shrivastava); General Hospital, Ernakulam (Balamuralikrishna, DMRT, J. Seios, MD, Sunithra Nair, MBBS); Government Medical College, Thrissur (K.L. Jayakumar, MD, Shehna A. Khader, DNB, M.B. Jayaraman, MD, V.R. Ajith Kumar, DMRT); Govt Arignar Anna Memorial Cancer Hospital & Research Institute, Kanchipuram (P. Anandhi, MD, C.T. Muthukumar, MBBS); HCG Bangalore Institute of Oncology, Bengaluru (BS Ajaikumar, MD, B.S. Ramesh, MD); HCG NMR Cancer Centre, Hubli (Sanjay Mishra, MD); Indira Gandhi Institute of Medical Sciences, Patna (Sangeeta Pankaj, MD, Rajesh Kumar Singh, MD); Indo-American Cancer Institute & Research Centre, Hyderabad (G. Dilip Kumar, DNB, Sujith Chayy Patanaik, MS, K. Kalpana Raghunath, MBBS, A. Santa, MD); International Cancer Centre, Neyyoor (Prarthana Roselil, MD, V.G. Sudhakaran, MD); Kidwai Memorial Institute of Oncology, Bengaluru (Ramachandra, MS, C. Ramesh, PhD); Mahavir Cancer Sansthan and Research Centre, Patna (Richa Chauhan, MD, Ravi Shankar Rajendra, MD, Rita Rani, DNB, Usha Singh, MD); Malabar Cancer Centre, Kannur (T. Bindhu, K. Ratheesan, Saina Sunilkumar, MD); Mandya Institute of Medical Sciences, Mandya (S. Shiva Kumar, MS, N. Venkatesh, MD); Max Super Specialty Hospital, New Delhi (Anil Kumar Anand, MD, Ramandeep Arora, MD); Max Super Specialty Hospital, Patparganj (Geeta Kadayaprath, MS, Meenu Walia, MD); Medanta Cancer Centre, Gurugaon (Tejinder Kataria, MD, Ashok Kumar Vaid, MD); MES Medical College & Hospital, Perinthalmanna (K.V. Gangadharan, MD, Mujeeb Rahman, MS); Mizoram State Cancer Institute (Civil Hospital), Aizwal (K. Lalfakzuala, MD, Lalhlupuii, MD, Doris Lallawmzuali, MD, Cindy Lalhanpuii, MD, Jeremy L. Pautu, MD, B. Zothankima, MD); Narayana Hrudayalaya Health City, Bengaluru (Alben Sigamani, MD); North East Cancer Hospital & Research Institute, Guwahati (Gazi Naseem Ahmed, MD, Imliwati Longkumer, D.K. Nath, MD); Post-graduate Institute of Medical Education and Research, Chandigarh UT (Sushmita Ghosal, MD, R. Kapoor, MD, Narender Kumar, MD); Pravara Rural Hospital & Rural Medical College, Loni (Vandana Jain, MD, Chaitali Wagmere, MD); Rajiv Gandhi Cancer Institute and Research Centre, New Delhi (Swarima Jaitley, Sudhir Rawal, MS); Regional Cancer Centre, Raipur (Pradeep Chandrakar, MD, Vivek Choudhary, MD); Regional Cancer Centre, Agartala (Dhritiman Datta, MD, Biswajit Debbamma, MD, Partha Sarathi Sutradhar, MD, Gautam Majumdar, MD); Regional Cancer Centre, Thiruvananthapuram (Preethi Sara George, PhD, Aleyamma Mathew, PhD); Regional Cancer Centre Kamala Nehru Memorial Hospital, Allahabad (Radha Ghosh, DNB, B. Paul Thaliath, MD); Regional Cancer Centre, Indira Gandhi Medical College, Shimla (Manish Gupta, MD, Sudarshan Sharma, MD); Regional Institute of Medical Sciences, Imphal (Indibor Singh, MD, Madhubala Devi, MD, Punyabati Devi, MD, Rajesh Singh Laishram, MD); RST Regional Cancer Hospital, Cancer Relief Society, Nagpur (Anjali Kolhe, MD, B.K. Sharma, MD); Rural Development Trust, Bathalapalle (N. Hariharanadha Sarma, MD); Sanjay Gandhi Post Graduate Institute of Medical Sciences, Lucknow (Rajesh Harsvardhan, MD, MD, Rakesh Pandey, MD, Punitalal, MD); SDM College of Dental Sciences and Hospital, Dharwad (Swetha Acharya, MS, Kaveri Hallikeri, MS); Shakuntala Memorial Hospital & Research Centre, Hubli (Vijay Gadagi, MD, Vinay Gadigi, MD); Sher-I-Kashmir Institute of Medical Sciences, Srinagar (Nazir Ahmad Khan, MD, Lone Mohammad Maqbool); St. Johns Medical Hospital, Bengaluru (Rakesh S Ramesh, DNB); Tata Medical Centre, Kolkata (Indranil Mallick, MD); Tata Memorial Hospital, Mumbai (B. Ganesh, PhD, R.A. Badwe, MS); The Gujarat Cancer & Research Institute, Ahmedabad (Shashank Pandya, MS, Anand Shah, MD); Vydehi Institute of Medical Sciences, Bengaluru (M.S. Ganesh, MS, Aruna E. Prasad, MD); JIPMER, Regional Cancer Centre, Puducherry UT (Biswajit Dubashi, MD, D. Kadambari, MD, Debasis Gochhait, MD, G.S. Sreenath, MD, K. Gunaseelan, K.T. Harichandrakumar, MD, Latha Chaturvedua, MD, Pampa Ch. Toi, MD, Pooja Sethi, MD, Prasanth Penumade, MD, B.H. Srinivas, MD, Sunil Kumar Saxena, MD); Government Medical College, Jammu (Dinesh Kumar, MD, Ashutosh Gupta, MD).